## Introduction to Quantum Computation, UPB Summer 2021, Assignment 1

To be completed by: Friday, April 23

## 1 Exercises

- 1. For complex number c = a + bi, recall that the *real* and *imaginary* parts of c are denoted Re(c) = a and Imag(c) = b.
  - (a) Prove that  $c + c^* = 2 \cdot \text{Re}(c)$ .
  - (b) Prove that  $cc^* = a^2 + b^2$ . How can we therefore rewrite |c| in terms of a and b?
  - (c) What is the polar form of  $c = \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}}i$ ? Use the fact that  $e^{i\theta} = \cos\theta + i\sin\theta$ .
  - (d) Draw  $c = \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}}i$  as a vector in the complex plane, ensuring to denote both the length of the vector and its angle with the x axis.
- 2. Prove that for any normalized vectors  $|\psi\rangle, |\phi\rangle \in \mathbb{C}^d$ ,

$$\| |\psi\rangle - |\phi\rangle \|_2 = \sqrt{2 - 2 \cdot \operatorname{Re}(\langle \psi | \phi \rangle)}.$$

Why does it not matter if we replace  $\langle \psi | \phi \rangle$  with  $\langle \phi | \psi \rangle$  in this equation?

3. Define

$$A = \left(\begin{array}{cc} a & b \\ c & d \end{array}\right).$$

- (a) What is  $\text{Tr}(A \cdot |1\rangle\langle 0|)$ ? (Hint: This can be computed quickly by using the cyclic property of the trace and the outer product representation of A. Do master this trick; it will be used repeatedly in the course and save you much time.)
- (b) Let  $|+\rangle = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$ . Use the same tricks as in part A, along with the fact that the trace is linear, to quickly evaluate

$$\operatorname{Tr}(A \cdot |+\rangle\langle +|).$$

- 4. (a) A general property of the outer product is that  $(|\psi\rangle\langle\phi|)^{\dagger} = |\phi\rangle\langle\psi|$ . Verify that this holds for the case where  $|\psi\rangle = |0\rangle$  and  $|\phi\rangle = |1\rangle$ . (Hint: Write out the full matrix corresponding to  $|0\rangle\langle1|$ .)
  - (b) Use Part (a) to prove that a normal matrix A satisfies  $A = A^{\dagger}$  if and only if all of A's eigenvalues are real. (Hint: Since A is normal, you can start by writing A in terms of its spectral decomposition. What does the condition  $A = A^{\dagger}$  enforce in terms of A's spectral decomposition?)